Ibuprofen (Advil/Motrin) Dosage

| | | Infant's Ibuprofen Concentrated Drops 50 mg (in each 1.25mL) | Children's Ibuprofen Suspension 100 mg (in each 5 mL) | Children's Soft Chewable Tablets 50 mg each | Junior Strength Chewable Tablets 100 mg each | Junior Strength Ibupofen Caplets 100 mg each |
|--------------|------------------|---|---|---|--|--|
| Weight | Age | Dropperful (Use dropper) | Teaspoon | Tablet | Tablet | Caplet |
| 12-17 lbs | 6-11 mos | 1.25 mL | | | | |
| 18-23 lbs | 12- 23 mos | 1.875 mL | | | | |
| 24-35 lbs | 2-3 yrs | 2.5 mL | 5 mL (1 tsp) | 2 | | |
| 36-47 lbs | 4-5 yrs | | 7.5 mL (1½ tsp) | 3 | | |
| 48-59 lbs | 6-8 yrs | | 12.5 mL (2 tsp) | 4 | 2 | 2 |
| 60-71 lbs | 9- 10 yrs | | 12.5 mL (2 ½ tsp) | 5 | 2 1/2 | 2 1/2 |
| 72-95 lbs | 11 yrs | | 15 mL (3 tsp) | 6 | 3 | 3 |

Please Note: One dose lasts 6-8 hours. Not to be used under 6 months.

- Milliliter is abbreviated as mL; 5mL equals one teaspoon (tsp).
- · Don't use household teaspoons, which can vary in size.
- · Aspirin should not be used in children to treat fever or pain.